

STUDY OF WATER QUALITY STATUS OF BORINA POND, CHARGAVAN, TEHSIL – RITHI, DISTRICT KATNI (M.P.), INDIA

Balmahendra Prajapati and Dr. Umesh Pandey
Department of Zoology, Govt. Science College, Rewa (M.P.), India

ABSTRACT: During last few years there has been an increasing trend to monitoring water quality of various fresh water bodies by regular measurement of their physico-chemical characteristics. Borina pond is situated in tehsil Rithi of district, Katni (M.P.) and lies at latitude 23°90" N and longitude 81°14" E. It is a perennial big pond and spread in about 15.40 hectares area. The depth of pond becomes about 15 meters in rainy season and mainly used for irrigation and fish culture. This study was carried out during November 2014 to October 2016. Four sampling stations were selected for the physico-chemical studies. Various physico-chemical parameters like water temperature, transparency, pH, dissolved oxygen, free carbon dioxide, biochemical oxygen demand, chemical oxygen demand, conductivity, total alkalinity, total solids, calcium hardness, magnesium hardness, total hardness, chloride contents, nitrate, phosphate and potassium contents were measured. From the study, it is inferred that the pond water is good for irrigation and fish culture. Most of the physico-chemical factors were found within the desirable limit of WHO except COD and total solids.

KEYWORDS: Water quality, Borina pond, physico-chemical parameters.

INTRODUCTION:-

Water is the most necessary component for the living being. Life on the earth is never possible without water. Shrivastava and Kanungo (2013) reported that about 70% of the available water in our country is polluted due to discharge of effluents from the industries, domestic waste, land and agricultural drainage. A healthy aquatic environment and production of sufficient fish food organisms in ponds are two factors of primary importance for successful pond cultural operations. To keep the aquatic habitat favorable for existence, physical and chemical factors will exercise their influence individually while the nutrient status of water play the most important role in governing the production of

plankton organisms. The object of the present study is to review and present a concise opinion for the optimum level of water quality parameters required for maximum fish productivity. Recently, many workers have studied the water quality of fresh water bodies of India i.e. Ansari et. al. (2012) on water quality of a Temple pond of Surat (Gujrat) India, Dixit et. al. (2015) on physico-chemical parameters of different pond water of Bilaspur district Chhattishgarh, India, Nagmani et. al. (2015) on water quality in five blocks of Bangalore, Sandhya and Benarjee (2016) on physico-chemical properties of some selected fresh water fish ponds in Warangal area of Telangana state, Kashyap (2016) on physico-chemical properties of water of Rewa (M.P.) and Younas et. al. (2017) on physico-chemical parameters of water and soils of three dams of district Korak (K.P.), Pakistan. The present study on water quality of Borina pond is aimed to assess the success of fish culture in relation to physiographic of the pond.

MATERIALS AND METHODS:-

The physico-chemical parameters of Borina pond were studied during two years of study period (November 2014 to October 2016) from four different sampling stations. Water samples were analyzed for some physico-chemical parameters. The yearly minimum and maximum mean values and mean \pm SD values are given table 1. The results obtained were compared with WHO and BIS standards. Water samples were collected in monthly interval from the surface of the pond at 9.30 am to 10.30 am in order to maintain uniformity. Samples were collected in one litre clean white polythene containers. Collected samples were brought to the laboratory and kept in the refrigerator for further analysis. Field parameters like temperature pH and transparency were determined at sampling site while other parameters like EC (\square mhos/cm), total solids (mg/l), TDS (mg/l), TSS (mg/l), total alkalinity (mg/l), calcium (mg/l), Mg (mg/l), total hardness (mg/l), chloride (mg/l), DO (mg/l), BOD (mg/l), COD (mg/l), phosphate (mg/l), chloride (mg/l), nitrate (mg/l) and potassium were analyzed in the laboratory by following the methodology of APHA (2005), Trivedi and Goel

(1984) and Wetzel and Likens (1991). Temperature, transparency, pH and conductivity were measured by using Celsius thermometer (0-110°C), Secchi disc, digital pH meter and conductivity meter respectively. Total solids, TDS and TSS were measured by volumetric analysis. Total alkalinity, total hardness, calcium, magnesium, chloride, free CO₂, DO, BOD, and COD were analyzed by titration method. Phosphate and Nitrate were analyzed by using UV-VIS spectrophotometer and potassium by flame photometer.

RESULT AND DISCUSSION:-

Minimum, maximum and mean±SD values of 17 selected physico-chemical parameters are given in Table 1. The water temperature was found to be ranged between 17.95°C to 26.68°C with mean±SD value of 22.38 ± 2.59°C during first year and between 18.08°C to 26.23°C with mean ± SD value of 22.51±2.35°C during second year of study period. The minimum values of water temperature were observed in the month of January and maximum in the month of May during study period. Shinde et. al. (2010) reported the range of water temperature between 17.0°C to 32.08°C in Harsool-Savangi dam, Maharashtra. Sandhya and Benarjee (2016) observed the water temperature between 26.8±2.90°C to 27.00±2.00°C in different ponds of Warangal area of Telangana. The ideal water temperature for biological activities of microorganisms is 20-25°C. Boyd (1998) reported the optimum range of water temperature between 25-32°C for tropical climate of a fish pond.

The mean values of transparency were recorded between 28.83 cm to 69.45 cm with mean±SD value of 49.99±14.12 cm during first year and between 28.45 cm to 69.75 cm with mean±SD value of 49.54±14.06 cm during second year of study period. The minimum transparency was observed in the month of July and maximum in the month of November during study period.

Boyd (1998) reported the optimum level of transparency between 30-40 cm. for fresh water aquaculture. USEPA (1974) reported that water transparency value less than 2.0 m is considered a eutrophic lake. Sawant and Chavan (2013) reported the transparency between 32.5 cm (July) to 62 cm (November) in Mahagaon reservoir, Gadhinglaj, Maharashtra. Querijero and Mercurio (2016) observed the highest transparency in the month of January and February in Taal lake, Batangas, Philippines. pH was recorded to varying from 7.28 to 8.78 with mean±SD value of 8.07±0.046 during first year and from 7.28±8.73 with mean value of 8.10±0.47 during second year of study period. The maximum pH was recorded in summer season and minimum in rainy season. Borina pond pH was within the desirable limits as set for protection of aquatic life by USEPA (6.5-9.0), BIS (6.5-

8.5) and ICMR (5.5-9.0). Sandhya and Benerjee (2016) reported the acceptable range of pH would be 6.5 to 9.0.

The recorded values of DO were ranged between 7.10 to 9.55 mg/l with mean±SD value of 8.07±0.46 mg/l and between 7.20 to 9.40 mg/l with mean±SD value of 8.05±0.75 mg/l during first and second years of study period respectively. The minimum values of DO were observed in summer season and maximum in winter season. WHO limits for DO was between 8-10 mg/l and BIS 6.0 mg/l. Banerjee and Babulal (1990) reported the pond water between 5.0 to 10.00 ppm of dissolved oxygen is ideal for fish culture. Sawant and Chavan (2013) reported the values of DO between 6.081 to 12.026 mg/l, minimum in the month of May and maximum in the month of December. Sandhya and Benarjee (2016) observed DO between 6.8±1.02 to 7.4±1.56 mg/l in Warangal area of Telangana.

During the study period, BOD was between 3.88 to 7.70 mg/l with mean ±SD value of 5.16±1.18 mg/l during first year and between 3.83 to 7.65 mg/l with mean± SD value of 5.14±1.22 mg/l during second year of study period. The higher values of BOD were recorded in summer season and lower in winter season. ICMR standard for BOD was 5.0 mg/l. WHO (1998) reported the minimum limit of pollution is indicated by BOD < 6mg/l. Boyd (1998) reported the optimum level of BOD < 10 mg/l for fresh water aquaculture. Sandhya and Benarjee (2016) reported the range of BOD between 1.89±0.074 to 3.14±0.015 mg/l in different ponds of Warangal area.

COD was found to be ranging between 16.13 to 23.65 mg/l with mean±SD value of 19.79±2.85 mg/l and between 16.38 to 22.55 mg/l with mean±SD value of 19.48±2.39 mg/l during first and second years of study period respectively. The maximum values of COD were recorded in summer season and minimum during winter season.

Boyd (1998) reported the optimum level of COD <50 mg/l for fresh water aquaculture. Kumar et. al. (2014) reported the COD between 10.85 mg/l (minimum in month of July) to 26.80 mg/l (maximum in February) in Yamuna river at Kalpi, Jalaun (U.P.).

During present study the minimum and maximum values of free CO₂ varied between 2.37 to 4.13 mg/l with mean±SD value of 3.30±0.54 mg/l during first year and between 2.32 to 4.13 mg/l with mean value of 3.25±0.55 mg/l during second year of study period. The higher values of free CO₂ were recorded during winter season and lower during summer seasons. Hynes (1970) reported 25 mg/l of free CO₂ as upper limit for fish culture. The values of free CO₂ in Borina Pond were found within tolerable limit for fish culture since it did not exceed 10 mg/l (APHA, 1998).

Table No. 1 - The magnitude of the mean values of Physico-Chemical factors of Borna Pond.

S.No.	Parameters	Units	Nov. 2014 to Oct. 2015			Nov. 2015 to Oct. 2016		
			Min.	Max.	Mean \pm SD value	Min.	Max.	Mean \pm SD value
1	Water temperature	°C	17.93	26.68	22.38 \pm 2.59	18.08	26.23	22.5 \pm 2.35
2	Transparency	cm.	28.83	69.45	49.99 \pm 14.12	28.45	69.75	49.54 \pm 14.06
3	pH	Units	7.28	8.78	8.07 \pm 0.46	7.28	8.73	8.10 \pm 0.47
4	DO	mg/l	7.10	9.55	8.07 \pm 0.77	7.20	9.40	8.05 \pm 0.75
5	BOD	mg/l	3.88	7.70	5.16 \pm 1.18	3.83	7.65	5.14 \pm 1.22
6	COD	mg/l	16.13	23.65	19.79 \pm 2.85	16.38	22.55	19.48 \pm 2.39
7	Free CO ₂	mg/l	2.37	4.13	3.30 \pm 0.54	2.32	4.13	3.25 \pm 0.55
8	Electrical conductivity	μ mhos/cm	160.50	327.25	229.42 \pm 49.40	161.25	319.25	228.23 \pm 48.62
9	Total alkalinity	mg/l	56.20	82.20	70.95 \pm 10.0	56.50	82.30	71.26 \pm 10.02
10	total solids	mg/l	423.25	961.25	607.13 \pm 173.44	421.75	951.25	605.60 \pm 172.00
11	Ca hardness	mg/l	80.25	153.58	107.56 \pm 24.68	79.00	150.33	106.79 \pm 24.30
12	Mg hardness	mg/l	13.28	25.45	18.03 \pm 3.77	13.10	25.28	18.14 \pm 3.84
13	Total hardness	mg/l	93.53	179.03	125.60 \pm 28.23	92.10	175.60	124.94 \pm 27.86
14	Chloride content	mg/l	15.40	27.15	20.87 \pm 3.30	15.95	27.23	20.96 \pm 3.13
15	Phosphate content	mg/l	0.30	0.56	0.41 \pm 0.08	0.28	0.54	0.39 \pm 0.09
16	Nitrate content	mg/l	0.19	0.45	0.30 \pm 0.08	0.21	0.43	0.31 \pm 0.07
17	Potassium content	mg/l	4.12	6.82	4.99 \pm 0.83	3.78	6.11	4.69 \pm 0.72

The values of electrical conductivity varied between 160.50 to 327.25 μ mhos/cm with mean \pm SD value of 229.42 \pm 49.40 μ mhos/cm during first year and between 161.25 to 319.25 μ mhos/cm with mean value of 228.23 \pm 48.62 μ mhos/cm during second year of study period. The values of electrical conductivity were recorded maximum in the month of May and minimum in the month of November during both study year. The BIS standard for electrical conductivity was 300 μ mhos/cm. Garg et. al. (2010) classified conductivity value greater than 500 μ mhos/cm as eutrophic. Sandhya and Benarjee (2016) reported the conductivity between 115.06 \pm 6.54 to 118.9 \pm 3.872 μ mhos/cm of the three ponds in Warangal area of Maharastra.

The mean values of alkalinity were varied between 56.20 to 82.20 mg/l with mean \pm SD value of 70.95 \pm 10.00 mg/l and between 56.50 to 82.30 mg/l with mean \pm SD value of 71.26 \pm 10.02 mg/l during first and second years of study period respectively. The minimum value of alkalinity was recorded in the month of September and maximum in the month of May during study period. Boyd (1998) reported the range of alkalinity between 50 to 300 mg/l for fish culture. The BIS standard for alkalinity was 600 mg/l. Chakravarty et. al. (2016) reported the alkalinity between 120 to 500 mg/l between different ponds of east

Godavari district. Andhra Pradesh. Thus, the value of alkalinity in Borina Pond has been found within the desirable limit of BIS standard.

The mean value of total solid varied between 423.25 to 961.25 mg/l with mean value of 607.13 \pm 173.44 mg/l during first year and between 421.75 to 951.25 mg/l with mean value of 605.60 \pm 172.00 mg/l during second year of study period. The minimum values of total solids were recorded in the month of January and maximum in the month of May during study period. Boyd (1998) reported the optimum water quality requirement for total solids <500 mg/l. The permissible limits for TDS as per WHO and BIS standard are 2000 mg/l and 2100 mg/l respectively (Shrivastva et. al., 2015). Younas et. al. (2017) reported that a sudden extreme changes in TDS could be killed aquatic life.

The minimum and maximum values of Ca hardness ranged between 80.25 to 153.58 mg/l with mean value of 107.56 \pm 24.68 during first year and between 79.00 to 150.33 mg/l with mean \pm SD value of 106.79 \pm 24.30 mg/l during second year of study period. The BIS standard for Ca hardness was 200 mg/l. Boyd (1998) reported the optimum level of calcium hardness between 75 to 150 mg/l for fresh water aquaculture.

The values of magnesium hardness varied between 13.28 to 25.45 mg/l with mean±SD value are 18.03±3.77 mg/l during first year and between 13.10 to 25.28 mg/l with mean±SD value of 18.14±3.84 mg/l during second year of study period. BIS standard for Mg was 100 mg/l.

The values of total hardness varied between 93.53 to 179.03 mg/l with mean+SD value of 125.60±28.23 mg/l in first year and between 92.10 to 175.60 mg/l with mean+SD value of 124.94±27.86 mg/l during second year of study period. The minimum value of total hardness were recorded in the month of January and maximum in the month of May during study period. The increase of hardness in summer month was due to the decrease in water level and increase in the rate of evaporation of high temperature. Sawant and Chavan (2013) reported the total hardness between 88.5 mg/l to 136.0 mg/l in Mahagaon reservoir (Maharashtra) with higher values in summer season.

The mean values of chloride contents varied between 15.40 to 27.15 mg/l with mean+SD value of 20.87±3.30 mg/l during first year and between 15.95 to 27.23 with mean+SD value of 20.96±3.13 mg/l during second year of study period. The chloride level of drinking water quality should be within 250 mg/l (WHO) and acceptable limit of BIS was 200 mg/l. Boyd (1998) reported the optimum level of chlorides between 31-50 mg/l for fresh water aquaculture. Kashyap (2016) reported the chloride content between 120 to 190 mg/l in various water samples of Rewa (M.P.). The chloride content of water of Borina pond was within desirable limit of WHO and BIS standards and suitable for fish culture.

The mean values of phosphate contents varied between 0.30 to 0.56 mg/l with mean value±SD value of 41±0.08 during first year and between 0.28 to 0.54 mg/l with mean value±SD value of 0.39±0.09 during second year of study period. The maximum values of phosphate were observed in rainy season and minimum in summer season. Abdar (2013) also observed the higher concentration of phosphorus during monsoon months and lower during summer months. The level of phosphate of Borina pond was found higher than standard of DENR (0.05<1.0 mg/l) recommended for class C water (aquaculture purpose) during the entire study period. Sawant and Chavan (2013) reported the range of phosphate contents between 0.0 mg/l to 0.225 mg/l in Mahagaon reservoir.

The mean values of nitrate content varied between 0.19 to 0.45 mg/l with mean± SD value of 0.30±0.08 mg/l during first year and between 0.21 to 0.43 mg/l with mean± SD value of 0.31±0.07 mg/l during second year of study period. Banerjee and Babulal (1990) mentioned that pond water having more than 1.00 ppm of nitrate is considered to be good for optimum production of fishes. WHO standard for nitrate was below 45 mg/l.

Chakravarty et. al. (2016) reported the range of nitrate between 3.36 mg/l to 6.40 mg/l in culture ponds of Andhra Pradesh. DENR (1990) reported 10 mg/l of nitrate for C class of water for growth of fish.

The minimum and maximum values of potassium content fluctuated between 4.12 to 6.82 mg/l with mean± SD value of 4.99±0.83 mg/l during first year and between 3.78 to 6.11 mg/l with mean± SD value of 4.69±0.72 during second year of study period. The maximum values of potassium content were observed in summer season and minimum in rainy season during study period. Boyd (1998) reported the optimum level of potassium contents between 0.5-10 mg/l for fresh water aquaculture. Sandhya and Benarjee (2016) observed the values of potassium between 1.73±0.001 to 2.1±0.042 mg/l in ponds of Warangal area.

REFERENCES:-

1. Abdar, M.R. (2013). "Physico-chemical characteristics and phytoplankton of. Morna lake, Shirala (M.S.) India," *Biolife.*, 2(1): 1-7.
2. Ansari, Ekhalak, Gadhia, Mohini and Surana, Ranjana (2012). Water quality of a Temple pond (Khajod) of Surat District, India. *Kolhapur, Proceeding of International Conference SWRDM-2012.* 27-30.
3. APHA (2005). American Public Health Association. Standard methods for examination of waters and wastewaters, Washington, DC, USA. 21st Edn.
4. Banerjee, R.K. and Babulal (1990). Role of soil and water in fish farming with special reference to primary production: In: *Technologies for inland fisheries development.* Ed. Sagunan V.V. and Utpal Bhaumik 1648.
5. BIS (1991). Bureau of Indian Standard Drinking Water Speciation, 1st revision, ISI: 10500.
6. Boyd, C.E. (1998). Water quality in ponds for aquaculture. Research and Development Series. Number 43. International Centre for Aquaculture and Aquatic Environments, Alabama Agricultural Experimental Station, Auburn University, Alabama :402.
7. Chakravarty, M.S., Ganesh, P.R.C., Amarnath, D., Shanthi Sudha, D. and Srinu Babu, T. (2016). Spatial variation of water quality parameters of shrimp (*Litopenaeus vannamei*) culture ponds at Narsapurapupeta, Kajuluru and Kaikavolu villages of East Godavari district, Andhra Pradesh. *International Journal of Fisheries and Aquatic Studies*, 4(4): 390-395.
8. DENR (Department of Environment and Natural Resources) (1990). Philippines Administrative Order No. 34 Series of 1990. Revised water usage and classification, Water Quality Criteria.

9. Dixit, A.K., Pandey, S.K., Mehta, R., Ahmad, N. and Pandey, Gunjan, J. (2015). Study of physico-chemical parameters of different pond water of Bilaspur District, Chhattishgarh, India. *Environmental Skeptics and Critics*, 4(3): 89-95.
10. Garg, Jivtesh, Nicola Bonini, Boris Kozinsky, and Nicola Marzari (2010). "Role of Disorder and Anharmonicity in the Thermal Conductivity of Silicon-Germanium Alloys: A First Principles Study." *Phys. Rev. Lett.* 106, 045901.
11. Hynes, H.B.N. (1970). *The ecology of running waters*. Univ. Toronto Press. Toronto. 555.
12. ICMR (1975). *Manual of standard of quality for drinking water supplies* (2nd ed.). Special Report Series No. 44. Indian Council of Medical Research. New Delhi: 27.
13. Kashyap, V.R. (2016). Physico-chemical analysis of various water samples of Rewa district (M.P.) India. *International Journal of Applied Research*; 2(1): 311-313.
14. Kumar, M., Singh, R., Chaurasia, S. and Khare, P. K. (2014). Physico-Chemical Examination of Loticwater of River Yamuna at Kalpi, District Jalaun, Uttar Pradesh, India. *J. Environ. Res. Develop. Journal of Environmental Research And Development*. 10(03): 529.
15. Nagamani, C., Saraswathi Devi, C. and Shalini, A. (2015). Physico-chemical analysis of water samples. *International Journal of Scientific and Engineering Research*, 6 (1) : 2149-2155.
16. Querijero, B.L. and Mercurio, A.L. (2016). Water quality in aquaculture and non-aquaculture sites in Taal lake, Batangas, Philippines, *Journal of Experimental Biology and Agricultural Sciences*, 4(1): 109-115.
17. Sandhya, K. and Benarjee, G. (2016). Physico-Chemical Properties of Some Selected Fresh Water Fish Ponds in Relation to Fish Production in Warangal Area. 6(4): 23-31.
18. Sawant, R. and Chavan, N.S. (2013). Water quality status of Mahagaon reservoir from Gadhinglaj Tahsil of Maharashtra. *International Journal of Science, Environment and Technology*. 2(6): 1196-1204.
19. Shinde, S.E., Pathan, T.S., Raut, K.S., Moore, P.R. and Sonawane, D.L. (2010). Seasonal variations in physico- chemical characteristics of Harsool-Savangi dam, District Aurangabad, India. *The Ecoscan* 4(1) 37-44.
20. Shrivastava, Asheesh, Tandon, Shalini, A. and Rakesh Kumar (2015). Water Quality Management Plan for Patalganga River for Drinking Purpose and Human Health Safety. *International Journal of Scientific Research in Environmental Sciences*, 3(2): 0071-0087.
21. Shrivastava, S. and Kanungo, V.K. (2013). Physico-Chemical Analysis of Pond Water of Surguja District, Chhattishgarh, India. *International Journal of Herbal Medicine*, 1(4): 35-43.
22. Trivedi, R.K. and Goel, P.K. (1984). *Chemical and Biological methods for water pollution status*. Environmental publication, Karad (India).
23. USEPA (United States Environmental Protection Agency)(1974). *Quality criteria for water*. Washington, D.C. USA.
24. Wetzel, R.G. and Likens G.E. (1991). *Limnological Analyses*. 2nd Edn. Springer-Verlag, New York.
25. WHO (World Health Organization)(1998). *Guidelines for Drinking Water Quality*. World Health Organization, Geneva.
26. Younas, S., Junaid, F., Gul, S., Rehman, H.U., Iqbal, N., Adan, K., Rauf, F., Ahmad, I., Najoom, S., Inayatullah, Sadia, H. and Usman, K. (2017). Physico-chemical parameters of water and soil of three Dams of district Karak, K.P., Pakistan. *Journal of Entomology and Zoology Studies*, 5(3): 317-322..