

ECOLOGICAL STUDY ON STERCULIA URENS: TRADITIONALLY IMPORTANT MEDICINAL TREE SATNA FOREST DIVISION (M.P.)

Dr. R.P.Sagar

Guest Lecturer Department of Botany
Govt. Kamla Nehru Girls College Balaghat (M.P.)

ABSTRACT:- *Sterculia urens* belonging to family Sterculiaceae. It is a medium sized deciduous tree met within Central Indian Forests. The plant parts are used in various ways for economic upliftments by the tribals. Non Timber Forest Produces (NTFPs) are basic needs and income generation opportunities for forest based tribal communities. Many multi-purpose tree species make broad their opportunity from only the single species. *Sterculia urens* one of the important NTFPs species having myriads of uses. Gum exudate of this tree is having large market value. This research is to highlight the importance of *S. urens* to the tribal peoples of Satna district (M.P.) India. *S. urens* can be very important species for tribal communities for food security, medicinal and employments generation for their sustainable livelihood development. *Sterculia urens* has a great economic importance. The medicinal and other importance of the plant has been noted from the mouth of tribal of Satna district. From very ancient it was used in various ways. The two important treatises on Ayurveda, Sushrita Samhita and Charak Samhita describes the activity of the plants. Thus the finally concluded from the study that the *Sterculia urens* Roxb. plant is of a great economic as well as Ecological significance. The species should be managed silviculturally for the better regeneration and maintenance in the nature.

KEYWORDS:- *Sterculia urens*, Kadaya, gum, NTFPs, tribal people Satna, medicinal value

INTRODUCTION:-

Sterculia urens Roxb. is moderate sized tree found in Central Indian Forest and other parts of India and also in the world. *Sterculia urens* belonging to family Sterculiaceae. Its Ayurvedic name is Kullu. It is a principal tree of hilly area of Central India and coromandel coast. It has also ecological importance in

Satpura hills. Indigenous tribes are entirely or partially reliant on forests for their subsistence call forests home and consider it a significant element of their social lives. Non-Timber Forest Products (NTFPs) are the sole natural resource that allows accessibility and subsistence to the world's poorest people, and they may play a critical role in the life of forest fringe tribal groups all over the world (Dolui *et. al.*, 2014). Non-timber forest products refer to all biological supplements acquired from natural forests for human use other than wood. Nearly 170 million people live in India's forests and surrounding regions, with more than half of them being tribal who rely on forest plants, particularly trees, for non-timber forest products. (Narayanan and colleagues, 2011).

NTFPs are also important for the livelihoods, food security, nutrition value well as for job creation of more than 80 million people (FAO, 2020). More than 500 plants NTFPs have been gathered by tribes, according to ethnobotanical experts (Rout *et al.*, 2010). Wild food plants, spices, honey, oils, fodder, gums, resins, gum-resins, colours, wax, lac, brooms, fibers, fuel wood, charcoal, fences, wildlife goods, and raw materials such as bamboo, cane, and other NTFPs are collected by tribal communities for livelihood, income generation or for their personnel uses (Bhattacharya and Hayat, 2004; Omkar *et al.*, 2012).

Forest resources have value to human beings. The forestry is an art and science of managing an devaluating forest resources with the object of producing goods and services on sustained basis of human society. Forests provide several products broadly classified as major and minor forest products which serve as back-bone in rural economy. In this research, we are discussing about economic and medicinal significance of *Sterculia urens* for local forest dwelling communities.

STUDY METHODS:-

An extensive literature survey was carried out for compilation of ecological and medicinal information on selected gum yielding woody plant species *Sterculia*

urens Roxb. The data were compiled on the traditional uses of various plant parts of these species along with industrial applications and various management practices as adopted by the local communities and the concerned state departments. The fieldwork was conducted in Satna forest division Madhya Pradesh of India. An ecological and medicinal survey was conducted eliciting information through personal interviews of villagers with the help of local assistants and also through direct and indirect observations made during the field surveys. Locals who practice traditional medical practices and local elder and knowledgeable people were interviewed for information on medicinal uses and availability of

selected gum yielding species in both the Indian states. The morphological characteristics of selected species were noted down during the survey period. Various gum harvesting practices from the selected species were also recorded.

RESULT AND DISCUSSION:-

Sterculia urens Roxb. (Family: Combretaceae) popularly known as ‘Karaya’ or ‘Kadaya’. *Sterculia urens* is commonly known as: Madhya Pradesh is Kullu, Gujarati: Kadayo/Kogdol, Konkani: Pandruk, Hindi: Kulu/Katira, English: Indian-Tragacanth

Taxonomic classification Kingdom	Plantae
Phylum	Tracheophyta
Class	Magnoliopsida
Order	Malvales
Family	Malvaceae
Genus	<i>Sterculia</i> L.
Species	<i>urens</i> Roxb

Morphological characteristics of *S. urens*

Characteristics	Details
Habit	Deciduous Tree
Height	Up to 15 m
DBH	2 m
Bark	Grey white or reddish (10-12 mm thickness)
Leaf	Palmately 5 lobed, 20-30 cm; alternate, crowded at the end of the branches
Flower	Greenish yellow, small in terminal pinnacles, bisexual
Fruit	Follicle 2-5 cm diameter, red, covered with stinging hairs
Seed	3-6 seeds, brown or black color, oblong.

Distribution

Sterculia urens are found at an elevation of 300-750 m. Temperature of its natural habitat varies from 40-48 °C to 0- 10 °C and rainfall varies from 750-1250 mm. It is generally found in hill slopes, ridges, rocky crevices, eroded slopes and survives in stony, rocky, shallow and ferruginous well drained soils (Sukhadiya *et al.*, 2019). *Sterculia urens* is naïve species in India, Laos, Myanmar, Sri Lanka, Thailand, Vietnam, etc. In India, it is found in tropical Himalayas, west and central India, throughout eastern and Western Ghats, including states; Andhra Pradesh, Assam, Maharashtra, Gujarat, Odisha, Rajasthan, Madhya Pradesh, Karnataka, Bihar, Chhattisgarh, Kerala, etc.

Importance of *Sterculia urens* Gum

Tribal peoples use this species as traditional medicine to cure various ailments. Almost all plant parts having medicinal values, however, the collection of gum serves as income and employment generation source since the *Sterculia urens* gum has significant importance in pharma, health care, food, cosmetics, waste management, paper-textile, composite fiber, and leather industries from a long period of time. Gum exudate of this tree is having large market value at globally (Dhiman *et al.*, 2019). *S. urens* gum is employed as a thickening ingredient, particularly in textile printing paste, and is also used commercially as a food additive. It is used as a pulp binder in the paper industry. It's also

employed in the pharmaceutical, cosmetic, and leather sectors as a tablet binder and gelling agent. (CSIR, 1976; Anderson and Wang, 1994; Nath and Nath, 2013). According to Persistence Market Research Pvt. Ltd., the global Karaya Gum market might reach US\$ 90.1 million by 2025. (www.prnewswire.com). During the forecast periods 2021- 2026, advances in global urbanization, greater usage of chemicals and materials in the sector, and a growth in international firms, retailers, and national supply chains might cause the Karaya Gum market to expand. (www.marketwatch.com).

Traditional uses of different parts of *Sterculia urens* by tribals-

Gum and resin: Bark yields gum which used in pharmaceutical preparations, tanneries, garbarti making, etc. (Omkar *et al.* 2012). *S. urens* tree gum exudes used in foodstuffs as emulsifiers, stabilizers and thickeners (Oak *et al.*, 2015). Gum of Karaya use to treat blisters, blood dysentery, dysentery, joint pain, stomach disorder, throat infection, tonic, jam, and confectionary. Emulsifier Thickener, Dental adhesive (Lujan-Medina *et al.* 2013), A small amount of gum and a mixture of CaCO₃ are used as antidote for snakebite and are useful both externally as well as orally, Extract of gum is applied locally to remove the spine from the skin, helps in treatment of Leucoderma and peptic ulcer, Regularize menstrual disorders (Jain *et al.* 2005), helps to remove blisters, joint pain, throat infections, thickening agent, especially in printing-paste for the textile industry. Being a good pulp binder it is used in paper industry. It is also used in pharmaceutical, cosmetic and leather industries (CSIR, 1976; Kala, 2016). It is used as tablet binder and gelling agent in pharmaceutical industries (Nath and Nath, 2013; Kala, 2016). The gum of *Sterculia urens* is also used commercially as food additives (Anderson and Wang, 1994; Kala, 2016). It is also having importance in gynecology (Kala, 2011), a fried mixture of resin and wheat flour/jaggery is given to women as a nutritious food, abdominal disorders (Dhiman *et al.*, 2019), regularize menstruation, burning sensation, bone fractures (Meena and Rao 2010) .

Bark: Mixture of *Sterculia stem* bark and *Haldinia cordifolia* with pepper, decoction given orally for 9 days to treat Leucorrhoea by tribes of Eastern ghat, Arunachal Pradesh (Ratnam and Raju, 2005), barks are also used for heals wound and throat infections (Oak *et al.*, 2015), rubbing the feet on the bark of *S. urens* can heal the cracks in the feet, where is half a cup of bark decoction taken once a day for 10– 12 days to clear the uterus by

Korku tribe of Amravati district, Maharashtra (Jagtap *et al.*, 2006), teaspoon of bark powder mixed with warm water once given helps in labor pains (Panduranga *et al.* 2011) and is taken orally to maintain menstrual cycle (Jain *et al.*, 2005), stem bark ground with turmeric and the filtrate is mildly heated and administered in 2 spoonfuls twice a day for 5 days to treat Rheumatoid Arthritis (Rao *et al.* 2016). Oligospermia is treated by soaking the bark of the twig for 10 days and drinking water on an empty stomach. (Murthy, 2012; Dhiman *et al.*, 2019).

Root: Root powder of *Sterculia urens* (Teklej) mixed with bark powder of *Ailanthus excelsa* and bark powder of *Madhuca longifolia* var. *latifolia*, leaf powder of *Vitex negundo* boiled in water which is used for bathing and 10 g of mixture taken with water to reduce body swelling uses by Korku tribe of Amravati district of Maharashtra (Jagtap *et al.* 2006).

Seed: Seeds use to treat wound healing and throat infections (Oak *et al.*, 2015).

Leaf: A leaf, gum and bark of Karaya are used for wound and throat infection (Oak *et al.*, 2015), Leaf juice applied externally to treat wound fractures and cracked skin leukemia by the tribals in the Kollihills, Eastern ghats, Tamilnadu (Vaidyanathan *et al.* 2013).

Tree branches: Cordage, facilitates child delivery, provides ease of delivery and is used as a tonic after childbirth (Kala, 2016; Kala, 2011). Leaves of *S. urens* are use as fodder for livestock of tribal communities (Omkar *et al.* 2012).

Other NTFPs uses of *S. urens* by tribals

Its bark is useful for making rope and rough cloth (Kala, 2011), yields fibre (Omkar *et al.* 2012). Its Seeds are edible (Omkar *et al.* 2012), eaten after roasting, it having nutritional value. Seeds and young tender roots are eaten in times of food crisis (Oak *et al.*, 2015) [20]. The seed oil is suitable for edible purposes and soap manufacturing. (Galla and Dubasi, 2010). The branch stalk is used as a toothbrush to relieve toothache (Padal and Vijayakumar, 2013). Its wood is used for making chandelier, pencil, wooden picture frames, packing, etc (Seth, 2003; Sivaraj *et al.* 2017).

CONCLUSION:-

Forest are valuable natural resources, the goods provided by forests are immense importance. Mans relation with plants is immortal; it is observed that man have always selected forests places for their habitat because they

satisfy a number of basic necessities of life. Almost every part of *S. urens*. are useful to cure various diseases. Other than the medicinal uses also useful for making rope, matches, clothes, toothbrush, pencil, wooden picture frames, packing etc. seed of *S. urens* also having high nutrition value and young tender roots and seed oil are edible. In nutshell, *S. urens* have multifarious uses and immense potential in tribal socio-economy in India. Thus the finally concluded from the study that the *Sterculia urens* Roxb. plant is of a great economic as well as Ecological significance. The species should be managed silviculturally for the better regeneration and maintenance in the nature.

REFERENCES:-

1. Anderson DMW, Wang WP. The tree exudate gums permitted in foodstuffs as emulsifiers, stabilisers and thickeners. *Chemistry and Industry of Forest Products*. 1994;14(2):73-83.
2. Arinathan V, Mohan VR, Britto A, Murugan C. Wild edibles used by Palliyars of the Western Ghats, Tamil Nadu. *Indian J Tradit. Know*. 2007;6:163-168.
3. Bhattacharya P, Hayat SF. Sustainable NTFP management for rural development: a case from Madhya Pradesh, India. *International Forestry Review*. 2004;6(2):161-168.
4. CSIR. The wealth of India. Publication and Information Directorate, Council of Scientific and Industrial Research, New Delhi, 1976, X.
5. Dhiman M, Singh A, Sharma MM. A review on *Sterculia urens* Roxb.: a boon to the livelihood for tribal people and industry. *Industrial Crops and Products*. 2019;130:341-351.
6. Dolui G, Chatterjee S, Chatterjee ND. The importance of non-timber forest products in tribal livelihood: A case study of santal community in Purulia District, West Bengal, 2014.
7. FAO, UNEP. The State of the World's Forests 2020. Forests, biodiversity and people. Rome, 2020. <https://doi.org/10.4060/ca8642en>
8. Galla NR, Dubasi GR. Chemical and functional characterization of Gum karaya (*Sterculia urens* L.) seed meal. *Food Hydrocolloid*. 2010;24:479-485.
9. Jagtap SD, Deokule SS, Bhosle SV. Some unique ethnomedicinal uses of plants used by the Korku tribe of Amravati district of Maharashtra, India. *J. Ethnopharmacol*. 2006;107:463-469. <https://doi.org/10.1016/j.jep.2006.04.002>.
10. Jain A, Katewa SS, Chaudhary BL, Galav P. Folk herbal medicines used in birth control and sexual diseases by tribals of southern Rajasthan, India. *J Ethnopharmacol*. 2004;90:171-177. <https://doi.org/10.1016/j.jep.2003.09.041>.
11. Jain A, Katewa SS, Galav PK, Sharma P. Medicinal plant diversity of Sitamata wildlife sanctuary, Rajasthan, India. *J Ethnopharmacol*. 2005;102:143-157.
12. Kala CP. Important gum yielding species *Anogeissus latifolia* (Roxb.) Bedd., *Boswellia serrata* Roxb. and *Sterculia urens* Roxb.: ethnobotany, population density and management. *Appl. Ecol. Env. Sci*. 2016;4:61-65. <https://doi.org/10.12691/aees-4-3-2>.
13. Kala CP. Indigenous uses and sustainable harvesting of trees by local people in Pachmarhi Biosphere Reserve of India. *Int. J. Med. Arom. Plants*, 2011, 153-161.
14. Lujan-Medina GA, Ventura J, Ceniceros ACL, Ascacio JA, Valdes DBV, Aguilar CN. Karaya gum: general topics and applications. *Macromolecules Indian J*. 2013;9:111-116.
15. Meena AK, Rao M. Folk herbal medicines used by the Meena community in Rajasthan. *Asian J Tradit. Med*. 2010;5:19-31.
16. Moe TS, Win HH, Hlaing TT, Lwin WW, Htet ZM, Mya KM. Evaluation of *in vitro* antioxidant, antiglycation and antimicrobial potential of indigenous Myanmar medicinal plants. *Journal of Integrative Medicine*. 2018;16(5):358- 366.
17. Murthy EN. Ethno medicinal plants used by gond of Adilabad district, Andhra Pradesh, India. *Int. J Pharm. Life Sci*. 2012;3:2034-2043.
18. Narayanan MR, Mithunlal S, Sujanalal P, Kumar NA, Sivadasan M, Alfarhan AH, *et al*. Ethnobotanically important trees and their uses by Kattunaikka tribe in Wayanad Wildlife Sanctuary, Kerala, India. *Journal of Medicinal Plants Research*. 2011;5(4):604-612.
19. Nath B, Nath LK. Evaluation of *Sterculia urens* gum as novel carrier for oral colon targeted drug delivery system. *Journal of Advanced Pharmaceutical Science and Technology*. 2013;1(2).
20. Oak G, Kurve P, Kurve S, Pejaver M. Ethnobotanical studies of edible plants used by tribal women of Thane District. *J Med. Plants Stud*. 2015;3:90-94.
21. Omkar K, Suthari S, Alluri S, Ragan A, Raju VS. Diversity of NTFPs and their utilization in Adilabad district of Andhra Pradesh, India. *Journal of Plant Studies*. 2012;1(1):33.

22. Padal SB, Vijayakumar Y. Traditional knowledge of Srikakulam District, Andhra Pradesh, India. *Int. J Innov. Res. Dev.* 2013;2:1310-1319.
23. Panduranga RM, Prasanthi S, Reddi ST. Medicinal plants in Folk medicine for Women's diseases in use by Konda Reddis. *Indian J Tradit. Know.* 2011;10:563-567.
24. Rao DS, Rao GMM, Murthy PP. Diversity and indigenous uses of some ethnomedicinal plants in Papikondalu wildlife sanctuary, Eastern Ghats of Andhra Pradesh, India. *Am. J Ethnomed.* 2016;3:06-25.
25. Ratnam VK, Raju VR. Folk medicine used for common women ailments Adivasis in the Eastern Ghats of Andhra Pradesh. *Indian J Tradit. Know.* 2005;4:267-270.
26. Rout SD, Panda SK, Mishra N, Panda T. Role of tribals in collection of commercial non-timber forest products in Mayurbhanj District, Orissa. *Studies of Tribes and Tribals.* 2010;8(1):21-25.
27. Sadia Raisa Lamia, Sumaiya Afra, Md Rezaul Hoque, Sajib Paul, Saifuzzaman Sumon. *Pharmacological*
28. Seth MK. Trees and their economic importance. *Bot. Rev.* 2003;69:321-376. [https://doi.org/10.1663/00068101\(2004\)069\[0321:TATEI\]2.0.CO;2](https://doi.org/10.1663/00068101(2004)069[0321:TATEI]2.0.CO;2).
29. Shukla A, Desai K, Modi N. *In vitro* antioxidant and antimicrobial potential of *Sterculia urens* Roxb. root extract and its bioactive phytoconstituents evaluation. *Future Journal of Pharmaceutical Sciences.* 2020;6(1):1-11.
30. Sivaraj N, Pandravada SR, Venkateswaran K, Dikshit N. Ethnic medicinal plant wealth of Eastern Ghats: status, knowledge systems and conservation strategies. *Int. J. Curr. Res. Biosci. Plant Biol.* 2017;4:83-101.
31. Sukhadiya M, Dholariya CA, Behera LK, Mehta AA, Patel DP, Gunaga RP. Commercial utilization and propagation of Gum Karaya: *Sterculia urens* Roxb. *MFP NEWS.* 2019;29(3):5-8
32. Vaidyanathan D, Senthilkumar MS, Basha MG. Studies on ethnomedicinal plants used by malayaltribals in Kolli hills of Eastern Ghats, Tamilnadu, India. *Asian J. Plant Sci. Res.* 2013;3:29-45.
33. www.marketwatch.com (Oct. 10, 2021)
34. www.prnewswire.com (Jan 16, 2018)
35. Yadav R, Yadav A. Phenology of selected woody species in a tropical dry deciduous forest in Rajasthan, India. *Trop. Ecol.* 2008;49:25-34.